

Friedberger Hochschulschriften

Ulrich Abel und Mircea Ivan

Asymptotic Expansion of the
Multivariate Bernstein Polynomials
on a Simplex

Friedberger Hochschulschriften Nr. 8

Das vorliegende Manuskript ist ein Preprint. Eine Arbeit mit den ausführlichen Beweisen wird in der mathematischen Fachzeitschrift Approximation Theory and its Applications erscheinen.

© Ulrich Abel

Friedberger Hochschulschriften
Herausgeber:
Die Dekane der Fachbereiche des Bereichs Friedberg der FH Gießen-Friedberg
Wilhelm-Leuschner-Straße 13, D-61169 Friedberg
<http://www.fh-friedberg.de>

Alle Rechte vorbehalten, Nachdruck, auch auszugsweise, nur mit schriftlicher Genehmigung und Quellenangabe.

Friedberg 2000
ISSN 1439-1112

Friedberger Hochschulschriften

Band 1: W. Hausmann

Das Nimspiel, der Assemblerbefehl XR und eine merkwürdige Art, zwei und zwei zusammenzählen

Band 2: U. Abel und M. Ivan

The Asymptotic Expansion for Approximation Operators of Favard-Szász Type

Band 3: C. Malerczyk

Visualisierungstechniken für den Sintflutalgorithmus

Band 4: M. Börgens, Th. Hemmerich und L. B. Rüssel

Use of Discriminant Analysis in Forecasting the Success of a Software Development Project

Band 5: W. Hausmann

On the two envelope paradox

Band 6: U. Abel

Asymptotic Approximation by Bernstein-Durrmeyer Operators and their Derivatives

Band 7: K. Behler

Hybrid Welding Technology (HWT) a flexible method for industrial applications

Band 8: U. Abel und M. Ivan

Asymptotic Expansion of the Multivariate Bernstein Polynomials on a Simplex

Asymptotic expansion of the multivariate Bernstein polynomials on a simplex

Ulrich Abel

Fachhochschule Giessen-Friedberg,

University of Applied Sciences, Fachbereich MND,

Wilhelm-Leuschner-Strasse 13, 61169 Friedberg, Germany

and

Mircea Ivan

Department of Mathematics,

Technical University of Cluj-Napoca,

Str. C. Daicoviciu 15, 3400 Cluj-Napoca, Romania

August 1, 2001

Abstract

In this note we study the local behaviour of the multi-variate Bernstein polynomials B_n on the d -dimensional simplex $S \subset \mathbb{R}^d$. For functions f admitting derivatives of sufficient high order in $\mathbf{x} \in S$ we derive the complete asymptotic expansion of $B_n f$ as n tends to infinity. All the coefficients of n^{-k} that only depend on f and \mathbf{x} are calculated explicitly. It turns out that combinatorial numbers play an important role. Our results generalize recent formulae due to R. Zhang.

1 Introduction

For each function f defined on $[0, 1]$, the classical Bernstein polynomials B_n ($n = 0, 1, 2, \dots$) are given by

$$B_n(f; x) = \sum_{\nu=0}^n \binom{n}{\nu} x^\nu (1-x)^{n-\nu} f\left(\frac{\nu}{n}\right) \quad (x \in [0, 1]). \quad (1)$$

Bernstein [8] already proved that, for $q \in \mathbb{N}$ and $f \in C^{2q}[0, 1]$, the univariate Bernstein polynomials satisfy the asymptotic relation

$$B_n(f; x) = f(x) + \sum_{k=1}^{2q} \frac{T_{n,s}(x)}{s! n^s} f^{(s)}(x) + o(n^{-q}) \quad (n \rightarrow \infty), \quad (2)$$

where

$$T_{n,s}(x) = \sum_{\nu=0}^n (\nu - nx)^s \binom{n}{\nu} x^\nu (1-x)^{n-\nu} \quad (s = 0, 1, 2, \dots).$$

In a very recent paper [19] R. Zhang presented a pointwise asymptotic expansion for Bernstein polynomials on a triangle in terms of the two-dimensional generalization of $T_{n,s}(x)$.

The drawback of Formula (2) and also of Zhang's bivariate formula is that the terms $T_{n,s}(x)$ contain the parameter n in a very implicit manner. The Bernstein polynomials B_n possess a complete asymptotic expansion of the form

$$B_n(f; x) \sim f(x) + \sum_{k=1}^{\infty} c_k(f; x) n^{-k} \quad (n \rightarrow \infty), \quad (3)$$

provided f admits derivatives of sufficiently high order at $x \in [0, 1]$. All coefficients $c_k(f; x)$ are independent of n . Formula (3) means that, for all $m = 1, 2, \dots$, there holds

$$B_n(f; x) = f(x) + \sum_{k=1}^m c_k(f; x) n^{-k} + o(n^{-m}) \quad (n \rightarrow \infty).$$

The aim of this paper is to derive a multi-dimensional version of Eq. (3) for multivariate Bernstein polynomials. All the coefficients of n^{-k} that only depend on f and x are calculated explicitly. It turns out that combinatorial numbers play an important role.

Let $\mathcal{S} \subset \mathbb{R}^d$ ($d \in \mathbb{N}$) be the simplex defined by

$$\mathcal{S} = \{ \mathbf{x} \in \mathbb{R}^d \mid x_i \geq 0 \ (i = 1, \dots, d), \ 1 - |\mathbf{x}| \geq 0 \}.$$

Throughout this paper, for $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$ and $\mathbf{k} = (k_1, \dots, k_d) \in \mathbb{N}_0^d$, we denote as usual

$$|\mathbf{x}| = \sum_{i=1}^d x_i, \quad \mathbf{x}^{\mathbf{k}} = \prod_{i=1}^d x_i^{k_i},$$

$$|\mathbf{k}| = \sum_{i=1}^d k_i, \quad \mathbf{k}! = \prod_{i=1}^d k_i!$$

and, for $\mathbf{k}, \mathbf{m} \in \mathbb{R}^d$, we write $\mathbf{k} \leq \mathbf{m}$ iff there holds $k_i \leq m_i$ ($i = 1, \dots, d$). For $n \in \mathbb{N}_0$, $\mathbf{k} \in \mathbb{N}_0^d$, we put $\binom{n}{\mathbf{k}} = n^{|\mathbf{k}|}/\mathbf{k}!$, where $x^{\mathbf{m}} = x(x-1)\cdots(x-m+1)$, $x^0 = 1$ denotes the falling factorial.

For each $n \in \mathbb{N}_0$ and $f : \mathcal{S} \rightarrow \mathbb{R}$, the multivariate Bernstein polynomial on the simplex \mathcal{S} is defined by

$$B_n(f; \mathbf{x}) = \sum_{|\mathbf{k}| \leq n} p_{n,\mathbf{k}}(\mathbf{x}) f\left(\frac{\mathbf{k}}{n}\right) \quad (\mathbf{x} \in \mathcal{S}), \quad (4)$$

where $p_{n,\mathbf{k}}(\mathbf{x}) = \binom{n}{\mathbf{k}} \mathbf{x}^{\mathbf{k}} (1 - |\mathbf{x}|)^{n-|\mathbf{k}|}$ and $\alpha \mathbf{x} = (\alpha x_1, \dots, \alpha x_d)$, for $\alpha \in \mathbb{R}$.

It is obvious that in the special case $d = 1$ we obtain the well-known univariate Bernstein polynomials (1). The case $d = 2$ are the Bernstein polynomials on a triangle which were studied by Zhang [19].

We mention that analogous results for the Bernstein–Kantorovich operators, the Meyer–König and Zeller operators and the operators of Butzer, Bleimann and Hahn can be found in [4, 1, 3, 2, 5]. Similar results on a certain positive linear operator can be found in [10, 7].

2 Main Results

Let $q \in \mathbb{N}$. For a fixed $\mathbf{x} = (x_1, \dots, x_d) \in \mathcal{S}$, let $K^{[q]}(\mathbf{x})$ be the class of all bounded functions $f : \mathcal{S} \rightarrow \mathbb{R}$ such that f and all its partial derivatives of order $\leq q$ are continuous in \mathbf{x} .

Theorem 1 (Complete asymptotic expansion for the operators B_n). *Let $q \in \mathbb{N}$, $\mathbf{x} \in \mathcal{S}$, and $f \in K^{[2q]}(\mathbf{x})$. The multivariate Bernstein polynomials on the simplex \mathcal{S} satisfy the asymptotic relation*

$$\begin{aligned} B_n(f; \mathbf{x}) &= f(\mathbf{x}) + \sum_{k=1}^q n^{-k} \sum_{\substack{\mathbf{s} \\ k < |\mathbf{s}| \leq 2k}} \frac{1}{\mathbf{s}!} \left(\frac{\partial^{|\mathbf{s}|}}{\partial x_1^{s_1} \cdots \partial x_d^{s_d}} f(\mathbf{x}) \right) \\ &\quad \sum_{\substack{\boldsymbol{\nu} \\ \boldsymbol{\nu} \leq \mathbf{s}}} a(k, \mathbf{s}, \boldsymbol{\nu}) \mathbf{x}^{\mathbf{s}-\boldsymbol{\nu}} + o(n^{-q}) \quad (n \rightarrow \infty), \end{aligned}$$

where the coefficients $a(k, \mathbf{s}, \boldsymbol{\nu})$ are given by

$$a(k, \mathbf{s}, \boldsymbol{\nu}) = \sum_{\substack{\mathbf{r} \\ \boldsymbol{\nu} \leq \mathbf{r} \leq \mathbf{s}, |\mathbf{r}| \geq k}} (-1)^{|\mathbf{s}| - |\mathbf{r}|} S(|\mathbf{r}| - |\boldsymbol{\nu}|, |\mathbf{r}| - k) \prod_{i=1}^d \left(\binom{s_i}{r_i} \sigma(r_i, r_i - \nu_i) \right). \quad (5)$$

Remark 1 . If $f \in K^{[\infty]}(\mathbf{x}) = \bigcap_{q=1}^{\infty} K^{[q]}(\mathbf{x})$, the multivariate Bernstein polynomials on the simplex \mathcal{S} possess the complete asymptotic expansion

$$B_n(f; \mathbf{x}) \sim f(\mathbf{x}) + \sum_{k=1}^{\infty} c_k(f; \mathbf{x}) n^{-k} \quad (n \rightarrow \infty),$$

where

$$c_k(f; \mathbf{x}) = \sum_{\substack{\mathbf{s} \\ k \leq |\mathbf{s}| \leq 2k}} \frac{1}{\mathbf{s}!} \left(\frac{\partial^{|\mathbf{s}|}}{\partial x_1^{s_1} \cdots \partial x_d^{s_d}} f(\mathbf{x}) \right) \sum_{\nu \leq \mathbf{s}} a(k, \mathbf{s}, \nu) \mathbf{x}^{\mathbf{s}-\nu}$$

and $a(k, \mathbf{s}, \nu)$ is as defined in (5).

The quantities $S(n, k)$ and $\sigma(n, k)$ denote the Stirling numbers of the first and second kind, respectively. Recall that the Stirling numbers are defined by the equations

$$x^n = \sum_{k=0}^n S(n, k) x^k \quad \text{resp.} \quad x^n = \sum_{k=0}^n \sigma(n, k) x^k \quad (n = 0, 1, \dots). \quad (6)$$

Furthermore we put $S(n, k) = \sigma(n, k) = 0$ if $k > n$.

Remark 2 . In the univariate case $d = 1$ we obtain the well-known formula

$$B_n(f; x) = f(x) + \sum_{k=1}^q n^{-k} \sum_{s=k}^{2k} \frac{1}{s!} f^{(s)}(x) \sum_{\nu=0}^s a(k, s, \nu) x^{s-\nu} + o(n^{-q})$$

as $n \rightarrow \infty$, where

$$a(k, s, \nu) = \sum_{r=\max\{\nu, k\}}^s (-1)^{s-r} \binom{s}{r} S(r-\nu, r-k) \sigma(r, r-\nu),$$

provided f is bounded on $[0, 1]$ and admits a derivative of order $2q$ at $x \in [0, 1]$ (cf. [2, Lemma 1]).

In the special case $q = 1$ Theorem 1 reveals the following Voronovskaja-type result.

Corollary 2 (Voronovskaja-theorem for the operators B_n). Let $\mathbf{x} \in \mathcal{S}$, and $f \in K^{[2]}(\mathbf{x})$. The multivariate Bernstein polynomials on the simplex \mathcal{S} satisfy the asymptotic relation

$$\lim_{n \rightarrow \infty} n(B_n(f; \mathbf{x}) - f(\mathbf{x})) = \frac{1}{2} \left(\sum_{i=1}^d x_i \frac{\partial^2 f(\mathbf{x})}{\partial x_i^2} - \sum_{i,j=1}^d x_i x_j \frac{\partial^2 f(\mathbf{x})}{\partial x_i \partial x_j} \right).$$

The classical theorem of Voronovskaja [18] is obtained in the special case $d = 1$.

For the convenience of the reader we list the explicit expressions for the initial terms of the asymptotic expansion. In order to simplify the notation we restrict ourselves to the case $d = 2$.

Let $(x, y) \in \mathcal{S}$, and $f \in K^{[6]}(x, y)$. The bivariate Bernstein polynomials on the simplex $\mathcal{S} \subset \mathbb{R}^2$ satisfy the asymptotic relation

$$\begin{aligned}
B_n f &= f + \frac{1}{2n} (x(1-x)f_{xx} - 2xyf_{xy} + y(1-y)f_{yy}) \\
&\quad + \frac{1}{6n^2} \left(\begin{array}{l} x(1-x)(1-2x)f_{xxx} + 3xy(2x-1)f_{xxy} \\ + 3xy(2y-1)f_{xyy} + y(1-y)(1-2y)f_{yyy} \end{array} \right) \\
&\quad + \frac{1}{8n^2} \left(\begin{array}{l} x^2(1-x)^2f_{x^4} - 4x^2y(1-x)f_{xy^3} \\ + 2xy(1-x-y+3xy)f_{x^2y^2} \\ - 4xy^2(1-y)f_{x^3y} + y^2(1-y)^2f_{y^4} \end{array} \right) \\
&\quad + \frac{1}{24n^3} \left(\begin{array}{l} x(1-x)(1-6x-6x^2)f_{x^4} - 4xy(1-6x-6x^2)f_{x^3y} \\ - 6xy(1-2x-2y+6xy)f_{x^2y^2} \\ - 4xy(1-6y-6y^2)f_{xy^3} + y(1-y)(1-6y-6y^2)f_{y^4} \end{array} \right) \\
&\quad + \frac{1}{12n^3} \left(\begin{array}{l} x^2(1-x)^2(1-2x)f_{x^5} - 5x^2y(1-x)(1-2x)f_{x^4y} \\ + xy(1-6x-y+15xy+5x^2-20x^2y)f_{x^3y^2} \\ + xy(1-x-6y+15xy+5y^2-20xy^2)f_{x^2y^3} \\ - 5xy^2(1-y)(1-2y)f_{xy^4} + y^2(1-y)^2(1-2y)f_{y^5} \end{array} \right) \\
&\quad + \frac{1}{48n^3} \left(\begin{array}{l} x^3(1-x)^3f_{x^6} - 6x^3y(1-x)^2f_{x^5y} \\ + 3x^2y(1-x)(1-x-y+5xy)f_{x^4y^2} \\ - 4x^2y^2(3-3x-3y+5xy)f_{x^3y^3} \\ + 3xy^2(1-y)(1-x-y+5xy)f_{x^2y^4} \\ - 6xy^3(1-y)^2f_{xy^5} + y^3(1-y)^3f_{y^6} \end{array} \right) \\
&\quad + o(n^{-3}) \quad (n \rightarrow \infty).
\end{aligned}$$

The corollary contains a result due to D. D. Stancu [17] (cf. [12, Eq. (5.87), p. 68]).

3 Auxiliary results

For each multi-index $\mathbf{r} = (r_1, \dots, r_d) \in \mathbb{N}_0^d$, we put $e_{\mathbf{r}}(\mathbf{x}) = \mathbf{x}^{\mathbf{r}}$.

Lemma 3. *For all $\mathbf{r} = (r_1, \dots, r_d) \in \mathbb{N}_0^d$, the moments of the Bernstein polynomials possess the representation*

$$B_n(e_{\mathbf{r}}; \mathbf{x}) = \sum_{k=0}^{|\mathbf{r}|} n^{-k} \sum_{\nu \leq \mathbf{r}, |\nu| \geq |\mathbf{r}| - k} \mathbf{x}^\nu S(|\nu|, |\mathbf{r}| - k) \prod_{i=1}^d \sigma(r_i, \nu_i) \quad (\mathbf{x} \in \mathcal{S}).$$

Proposition 4 . For all $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{N}_0^d$, the central moments of the Bernstein polynomials possess the representation

$$B_n((\cdot - \mathbf{x})^\mathbf{s}; \mathbf{x}) = \sum_{k=0}^{|\mathbf{s}|} n^{-k} \sum_{\nu \leq \mathbf{s}} a(k, \mathbf{s}, \nu) \mathbf{x}^{\mathbf{s}-\nu} \quad (\mathbf{x} \in \mathcal{S}), \quad (7)$$

where the coefficients $a(k, \mathbf{s}, \nu)$ are given by Eq. (5).

In order to show Theorem 1 we use a general approximation theorem for positive linear operators [6].

Lemma 5 . Let $q \in \mathbb{N}$ and $\mathbf{x} \in \mathcal{S}$. Moreover, let $L_n : K^{[2q]}(\mathbf{x}) \rightarrow C(\mathcal{S})$ be a sequence of positive linear operators. If, for $k = 2q$ and $k = 2q + 2$,

$$L_n(\|\cdot - \mathbf{x}\|_2^k; \mathbf{x}) = \mathcal{O}(n^{-\lfloor (k+1)/2 \rfloor}) \quad (n \rightarrow \infty), \quad (8)$$

then we have, for each $f \in K^{[2q]}(\mathbf{x})$,

$$L_n(f; \mathbf{x}) = \sum_{\substack{\mathbf{s} \\ |\mathbf{s}| \leq 2q}} \frac{1}{\mathbf{s}!} \left(\frac{\partial^{|\mathbf{s}|} f(\mathbf{x})}{\partial x_1^{s_1} \cdots \partial x_d^{s_d}} \right) L_n((\cdot - \mathbf{x})^\mathbf{s}; \mathbf{x}) + o(n^{-q}) \quad (n \rightarrow \infty).$$

The special case $d = 1$ is due to Sikkema [15, Theorems 1 and 2].

Remark 3 . For $r = 0, 1, 2, \dots$, we have

$$\|\mathbf{t} - \mathbf{x}\|_2^{2r} = \left(\sum_{i=1}^d (t_i - x_i)^2 \right)^r = \sum_{|\mathbf{s}|=r} \binom{r}{\mathbf{s}} (\mathbf{t} - \mathbf{x})^{2\mathbf{s}}.$$

In order to apply Lemma 5, we have to check whether the Bernstein polynomials satisfy condition (8). By Remark 3, we have to show that

$$L_n((\cdot - \mathbf{x})^{2\mathbf{s}}; \mathbf{x}) = \mathcal{O}(n^{-|\mathbf{s}|}) \quad (n \rightarrow \infty),$$

for all $\mathbf{s} \in \mathbb{N}_0^d$. In the following Lemma we shall prove a slightly more general result.

Lemma 6 . For each $\mathbf{x} \in \mathcal{S}$ and all $\mathbf{s} \in \mathbb{N}_0^d$, the central moments of the Bernstein polynomials satisfy the estimation

$$B_n((\cdot - \mathbf{x})^\mathbf{s}; \mathbf{x}) = \mathcal{O}(n^{-\lfloor (|\mathbf{s}|+1)/2 \rfloor}) \quad (n \rightarrow \infty).$$

For the proof we shall make use of the following well-known representations for the Stirling numbers of the first kind

$$S(n, n-k) = \sum_{j=k}^{2k} \binom{n}{j} S_2(j, j-k) \quad (k = 0, 1, \dots, n; n = 0, 1, 2, \dots) \quad (9)$$

resp. second kind

$$\sigma(n, n - k) = \sum_{j=k}^{2k} \binom{n}{j} \sigma_2(j, j - k) \quad (k = 0, 1, \dots, n; n = 0, 1, 2, \dots) \quad (10)$$

(see [9, p. 227], cf. [11, pp. 149–153]). The quantities $S_2(n, k)$, $\sigma_2(n, k)$ denote the associated Stirling numbers of the first resp. second kind. Recall that the associated Stirling numbers are defined by their double generating function

$$\sum_{n,k \geq 0} S_2(n, k) t^n u^k / n! = e^{-tu} (1+t)^u$$

resp.

$$\sum_{n,k \geq 0} \sigma_2(n, k) t^n u^k / n! = \exp(u(e^t - 1 - t)).$$

(see [9, p. 295 and p. 222]).

REFERENCES

- [1] U. Abel, *The moments for the Meyer-König and Zeller operators*, J. Approx. Theory **82** (1995), 352–361.
- [2] U. Abel, *On the asymptotic approximation with operators of Bleimann, Butzer and Hahn*, Indag. Math., (N.S.), **7**(1) (1996), 1–9.
- [3] U. Abel, *The complete asymptotic expansion for Meyer-König and Zeller operators*, J. Math. Anal. Appl. **208** (1997), 109–119.
- [4] U. Abel, *Asymptotic approximation with Kantorovich polynomials*, Approx. Theory and Appl. **14:3** (1998), 106–116.
- [5] U. Abel, *On the asymptotic approximation with bivariate operators of Bleimann, Butzer and Hahn*, J. Approx. Theory **97** (1999), 181–198.
- [6] U. Abel, *On the asymptotic approximation with bivariate Meyer-König and Zeller operators*, submitted.
- [7] U. Abel and M. Ivan, *Asymptotic approximation with a sequence of positive linear operators*, to appear in J. Comp. Math. Appl.
- [8] S. N. Bernstein, *Complément à l'article de E. Voronowskaja*, Dokl. Akad. Nauk SSSR **4** (1932), 86–92.
- [9] L. Comtet, “Advanced Combinatorics”, Reidel Publishing Comp., Dordrecht, 1974.
- [10] M. Ivan and I. Raşa, *A sequence of positive linear operators*, Rev. Anal. Numér. Théor. Approx. **24**(1-2) (1995), 159–164.

- [11] C. Jordan, “*Calculus of finite differences*”, Chelsea, New York, 1965.
- [12] G. Meinardus, “*Approximation von Funktionen und ihre numerische Behandlung*”, Springer, Berlin, 1964.
- [13] J. Riordan, “*Combinatorial identities*”, Wiley, New York, 1968.
- [14] P. C. Sikkema, *On some linear positive operators*, Indag. Math. **32** (1970), 327–337.
- [15] P. C. Sikkema, *On the asymptotic approximation with operators of Meyer-König and Zeller*, Indag. Math. **32** (1970), 428–440.
- [16] D. D. Stancu, *Some Bernstein polynomials in two variables and their applications*, Dokl. Akad. Nauk. SSSR **134**(1) (1960), 48–51.
- [17] D. D. Stancu, *Some Bernstein polynomials in two variables and their applications*, Soviet. Math. **1** (1960), 1025–1028.
- [18] E. V. Voronovskaja, *Determination de la forme asymptotique d'approximation des functions par les polynomes de M. Bernstein*, Dokl. Akad. Nauk. SSSR (A) (1932), 79–85.
- [19] R. Zhang, *An asymptotic expansion formula for Bernstein polynomials on a triangle*, Approx. Theory and Appl. **14:1** (1998), 49–56.